

Embargo Copy - Not for Distribution




PEDIATRICS

Sudden Unexpected Infant Deaths: 2015–2020

Carrie K. Shapiro-Mendoza, PhD, MPH, Kate R. Woodworth, MD, MPH, Carri R. Cottengim, MS, Alexa B. Erck Lambert, MPH, Elizabeth M. Harvey, PhD, MPH, Michael Monsour, PhD, Sharyn E. Parks, PhD, MPH, Wanda D. Barfield, MD, MPH

(doi: 10.1542/peds.2022-058820)

Embargo Release Date: Monday, March 13, 2023 - 12:01 am (ET)



Embargo Policy:

Information in this article is embargoed for release until the date indicated above. Interviews may be conducted prior to the embargo release date, but nothing may be aired or published.

If you are a media representative and have questions about the embargo, upcoming press events, or other matters, please contact AAP Communications staff at 630-626-6177, or via e-mail at commun@aap.org

Sudden Unexpected Infant Deaths: 2015–2020

Carrie K. Shapiro-Mendoza, PhD, MPH,^a Kate R. Woodworth, MD, MPH,^b Carri R. Cottengim, MS,^a
Alexa B. Erck Lambert, MPH,^c Elizabeth M. Harvey, PhD, MPH,^a Michael Monsour, PhD,^a Sharyn E. Parks, PhD, MPH,^a
Wanda D. Barfield, MD, MPH^a

abstract

OBJECTIVE: Although the US infant mortality rate reached a record low in 2020, the sudden infant death syndrome (SIDS) rate increased from 2019. To understand if the increase was related to changing death certification practices or the coronavirus disease 2019 (COVID-19) pandemic, we examined sudden unexpected infant death (SUID) rates as a group, by cause, and by race and ethnicity.

METHODS: We estimated SUID rates during 2015 to 2020 using US period-linked birth and death data. SUID included SIDS, unknown cause, and accidental suffocation and strangulation in bed. We examined changes in rates from 2019 to 2020 and assessed linear trends during prepandemic (2015–2019) using weighted least squares regression. We also assessed race and ethnicity trends and quantified COVID-19–related SUID.

RESULTS: Although the SIDS rate increased significantly from 2019 to 2020 ($P < .001$), the overall SUID rate did not ($P = .24$). The increased SIDS rate followed a declining linear trend in SIDS during 2015 to 2019 ($P < .001$). Other SUID causes did not change significantly. Our race and ethnicity analysis showed SUID rates increased significantly for non-Hispanic Black infants from 2019 to 2020, widening the disparities between these two groups during 2017 to 2019. In 2020, <10 of the 3328 SUID had a COVID-19 code.

CONCLUSIONS: Diagnostic shifting likely explained the increased SIDS rate in 2020. Why the SUID rate increased for non-Hispanic Black infants is unknown, but warrants continued monitoring. Interventions are needed to address persistent racial and ethnic disparities in SUID.

Full article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2022-058820

^aDivision of Reproductive Health and ^bCOVID-19 Response, Centers for Disease Control and Prevention, Atlanta, Georgia; and ^cDB Consulting Group, Inc, Silver Spring, Maryland

Dr Shapiro-Mendoza conceptualized and designed the study, carried out the initial analyses, drafted the initial manuscript, and reviewed and revised the manuscript; Dr Monsour carried out the initial analyses, assisted in interpreting study findings, and reviewed and revised the manuscript; Drs Barfield, Harvey, Parks, and Woodworth, Ms Cottengim, and Ms Erck Lambert assisted in interpreting study findings and critically reviewed and revised the manuscript for important intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2022-058820>

Accepted for publication Jan 3, 2023

Address correspondence to Carrie K. Shapiro-Mendoza, PhD, MPH, Division of Reproductive Health, Centers for Disease Control and Prevention, 4770 Buford Hwy NE, Mail Stop 107-2, Chamblee, GA 30341. E-mail: ayn9@cdc.gov

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2023 by the American Academy of Pediatrics

WHAT'S KNOWN ON THIS SUBJECT: United States infant mortality reached a record low in 2020. The sudden infant death syndrome (SIDS) rate increased 15% from 2019 to 2020. Because of diagnostic shifting, changes in SIDS rates must be considered with other sudden unexpected infant deaths.

WHAT THIS STUDY ADDS: Increased SIDS rates in 2020 was likely attributed to diagnostic shifting. Evidence does not support direct or indirect effects of the coronavirus disease 2019 pandemic on increased rates of sudden unexpected infant death, except for non-Hispanic Black infants.

To cite: Shapiro-Mendoza CK, Woodworth KR, Cottengim CR, et al. Sudden Unexpected Infant Deaths: 2015–2020. *Pediatrics*. 2023;151(4):e2022058820

In 2020, the United States achieved a record low infant mortality rate: 541.9 infant deaths per 100 000 live births.¹ Concurrently, the sudden infant death syndrome (SIDS) rate increased 15% from 2019 to 2020, moving from the fourth to the third leading cause of infant death.¹ It is unknown why the SIDS rate increased, but potential reasons may be related to the coronavirus disease 2019 (COVID-19) pandemic or changing death certification practices.

The COVID-19 pandemic may have affected sudden unexpected infant deaths (SUID) in several ways. Increased pressures on families, caregivers, and communities during the pandemic may have limited opportunities for clinical follow-up after birth (eg, early postpartum discharges and timely well-baby visits after birth hospitalization) and access to safety-net supports (eg, Women, Infants, and Children Supplemental Nutrition Program; Healthy Start).² Limited access to these encounters and services might have increased the risk of SIDS and other SUIDs. Clinical follow-up and support services are times when safe sleep messages are communicated, smoking cessation is encouraged, immunizations are administered, and breastfeeding counseling is given, all of which are protective factors against SIDS and SUID.^{3–5} Furthermore, pandemic-related losses of or changes in child care arrangements may also have led to an increased SIDS or SUID risk due to unaccustomed infant sleep positioning and other unsafe sleep practices.^{6,7} Monitoring changes in SUID rates before and during the COVID-19 pandemic can provide insight to understand if pandemic factors affected SIDS and SUID rates.

The increased SIDS rate previously reported in 2020¹ may also be related to diagnostic shifting as

opposed to indirect and direct effects of the COVID-19 pandemic. To understand reasons for the increase, it is important to examine SUID rates as a group in addition to cause-specific rates, and not solely SIDS in isolation. The reported 2020 increase was surprising because in recent years, the term SIDS had fallen out of favor among some medical examiners and coroners; SIDS is used less frequently in favor of other SUID causes, namely unknown cause and accidental suffocation and strangulation in bed (ASSB).^{8–12} This diagnostic shifting is well recognized. Until we have standardized reporting practices, monitoring SUID as a group, and not solely monitoring SIDS, is important given the variability in reporting practices and resultant diagnostic shift in the classification of SUID and its subtypes.^{10–13}

We explore US mortality data to better understand the epidemiology of SIDS and SUID during 2015 to 2020 and what may have led to the increased SIDS rate from 2019 to 2020. First, we describe overall and cause-specific SUID rates by year and assess trends during 2015 to 2020. Second, we explore changes in SUID rates by race and ethnicity. Because SUID rates may have been affected by COVID-19 or efforts to mitigate COVID-19 exposure, we quantify the number of SUID cases in 2020 that also had COVID-19 listed as a contributing cause of death. Having a complete understanding of changing SUID rates and disparities can help program planners and policymakers adapt and promptly implement risk reduction strategies and interventions for infants at highest risk.

METHODS

We used US period-linked birth and infant death files for 2015 to 2020.¹⁴ This period file was selected

to examine changes in SUID before (2015–2019) and during (2020) the COVID-19 pandemic; 2020 is the last year final national SUID data currently available. We calculated overall and cause-specific SUID rates per 100 000 live births along with 95 confidence intervals. In the period-linked file, infants (<365 days) who died in the United States in a given calendar year are linked back with the corresponding birth certificate.¹⁵ Analyses were restricted to infants born to US residents from the 50 states and District of Columbia. SUID included 3 underlying cause-of-death classifications^{16,17} coded per the International Classification of Diseases, 10th Revision (ICD-10): SIDS (R95), unknown cause (R99), and ASSB (W75).¹⁸ The ICD-10 is used for official reporting of mortality statistics in the United States and other nations.¹⁵ By law, the official cause of death for SUID cases must be determined and reported by the medicolegal death investigation system (ie, medical examiner and coroners).

We used the overall SUID rate (as opposed to cause-specific rates) to assess rates by race and ethnicity. Using this SUID grouping allowed for consistent comparisons unaffected by death certifier variations in reporting.⁸ In addition, the overall SUID rate is less likely to be affected by random variation in small death counts that might occur with some racial and ethnic groups.⁸ SUID rates by race and ethnicity were calculated as the number of SUID per 100 000 live births in the specified race and ethnic group in a given year.

We report single race and Hispanic origin groups per the 1997 Office of Management and Budget (OMB) standards¹⁹: non-Hispanic White, non-Hispanic Black, non-Hispanic American Indian or Alaska Native, non-Hispanic Asian, non-Hispanic

Native Hawaiian or Other Pacific Islander, and Hispanic. The 2017 period-linked file was the first year for which all 50 states and Washington DC had birth certificate data on race and Hispanic-origin categories by using the 1997 OMB standards. For consistency, analyses of single race and Hispanic origin groups were restricted to 2017 to 2020. Data before 2017 did not follow the 1997 standards and used bridged-race categories (not available in the 2020 data).¹⁵

COVID-19 status was estimated by using the ICD-10 code, U07.1.²⁰ This code should be reported when COVID-19 is assumed to have caused or contributed to death. The code was implemented March 24, 2020, in the United States and applied retroactively to cover the entire year.

To quantify the magnitude and direction of changes in SUID rates from 2019 to 2020, we calculated percent change^{15,21}; significance levels were adjusted for multiple comparisons by using the Bonferroni method.²² We then compared the direction of rate changes from 2019 to 2020 to prepandemic trends (2015–2019 and 2017–2019 for race and ethnicity). Prepandemic trends were measured using a weighted least squares regression of the mortality rates over time (years). Finding an increasing or decreasing rate from 2019 to 2020 inconsistent with the prepandemic rates and trends may provide preliminary evidence of a changing pattern. To guide our statistical testing and analysis for linear trends, we followed guidance by National Center for Health Statistics.²² *P* values <.05 were considered statistically significant.

Line graphs were plotted to show SUID changes over time by cause of death and race-ethnicity groups.

Analyses were conducted by using SAS version 9.4 (SAS Institute, Inc, Cary, NC). SUID rates based on <20 deaths were flagged unreliable and suppressed. This analysis did not require institutional review board review because it used deidentified public-use data sets without identifiable private information.

RESULTS

SUID Mortality

Of the 19 446 infant deaths in 2020, 3328 (17.1%) were classified as SUID (combination of R95, R99, and W75). The SUID rate per 100 000 live births increased from 89.5 in 2019 to 92.1 in 2020, but this increase was not significant (Table 1 and Fig 1). In addition, a significant linear trend was not observed for 2015 to 2019 (Fig 1).

In our analyses of specific causes of SUID, we observed that the SIDS (R95) rate was higher than rates for unknown cause (R99) and ASSB (W75) for all study years (Table 1 and Fig 1). Unlike the SUID rate, the SIDS rate increased significantly from 33.3 in 2019 to 38.2 in 2020, and a significant decreasing linear trend in the SIDS rate was observed during 2015 to 2019 (<.001) (Fig 1). For unknown cause and ASSB, the rates did not change significantly from 2019 to 2020 (30.8 vs 28.9 and 25.4 vs 25.0, respectively), and no significant trend was observed during 2015 to 2019.

The proportion of SUID by each cause fluctuated during 2015 to 2020 (Fig 2). Among SUID in 2020, 41% were classified as SIDS (range 37%–43% during 2015–2019). For all years, the percentage of SUID classified as SIDS was higher than deaths classified as cause unknown (range 31%–37%) and ASSB (range 24%–28%).

Race and Ethnicity

In 2020, the SUID rate per 100 000 live births was highest among non-Hispanic Black (214.0) and non-Hispanic American Indian or Alaskan Native (AIAN) infants (205.1), nearly 3 times the rate for non-Hispanic White infants (75.6) (Table 1 and Fig 3). Non-Hispanic Asian (23.3) and Hispanic infants (59.0) had the lowest rates. The 2020 SUID rate for non-Hispanic Black infants (214.0) was higher than any time during 2017 to 2019 (range 187.3–192.1). The 2020 SUID rates for non-Hispanic AIAN and non-Hispanic White infants were lower than any time during 2017 to 2019. A significant decreasing trend was observed for non-Hispanic White infants during 2017 to 2019 (*P* = .02). For Hispanic and non-Hispanic Asian infants, SUID rates generally increased, but no significant trends were observed. The SUID ratio for non-Hispanic Black infants compared to non-Hispanic White infants increased yearly during 2017 to 2020 from 2.2 in 2017 to 2.8 in 2020.

COVID-19

Among the 19 446 infant deaths in 2020, 35 deaths (0.2%) had COVID-19 listed as an underlying cause of death. Among the 3328 SUID in 2020, <10 deaths had a COVID-19 code indicating an additional (not an underlying) cause of death.

DISCUSSION

Although the SIDS rate increased significantly from 2019 to 2020, a significant increase in the overall SUID and other cause-specific SUID rates were not found. Interestingly, the increase in SIDS was unique to 2020 and contrasted to the declining linear trend in SIDS rates during 2015 to 2019. The lack of increasing SUID rates together with the small number of SUID that also had a COVID-19 cause-of-death code makes it unlikely that increased

TABLE 1 SUID Mortality Rates by Cause (2015–2020) and Race-Ethnicity (2017–2020), United States

	Number of SUID						SUID Rate per 100 000 Live Births ^a (95% Confidence Interval)					
	2015	2016	2017	2018	2019	2020	2015	2016	2017	2018	2019	2020
Total SUID ^b	3661	3585	3577	3428	3354	3328	92.0 (89.0–95.0)	90.9 (87.9–93.8)	92.8 (89.7–95.8)	90.4 (87.4–93.4)	89.5 (86.5–92.5)	92.1 (89.0–95.2)
Cause-specific												
SIDS	1563	1494	1357	1327	1247	1380	39.3 (37.3–41.2)	37.9 (35.9–39.8)	35.2 (33.3–37.1)	35.0 (33.1–36.9)	33.3 (31.4–35.1)	38.2 (36.2–40.2)
Unknown	1180	1234	1275	1266	1155	1046	29.7 (28.0–31.4)	31.3 (29.5–33.0)	33.1 (31.3–34.9)	33.4 (31.5–35.2)	30.8 (29.0–32.6)	28.9 (27.2–30.7)
ASSB	918	857	945	835	952	902	23.1 (21.6–24.6)	21.7 (20.3–23.2)	24.5 (22.9–26.1)	22.0 (20.5–23.5)	25.4 (23.8–27.0)	25.0 (23.3–26.6)
Race and ethnicity ^c												
Hispanic ^d	—	—	502	487	523	511	—	—	55.9 (51.0–60.7)	55.0 (50.1–59.8)	59.0 (53.9–64.1)	59.0 (53.8–64.1)
NH AIAN	—	—	69	61	59	55	—	—	230.3 (179.2–291.5)	209.7 (160.4–269.3)	207.4 (157.9–267.5)	205.1 (154.5–267.0)
NH Asian	—	—	46	65	52	51	—	—	18.5 (13.5–24.6)	27.0 (20.8–34.4)	21.8 (16.3–28.6)	23.3 (17.3–30.6)
NH Black	—	—	1077	1034	1041	1134	—	—	192.1 (180.6–203.6)	187.3 (175.9–198.7)	189.9 (178.4–201.5)	214.0 (201.6–226.5)
NH NHOPI	—	—	13	25	14	11	—	—	e	e	e	e
NH White	—	—	1703	1584	1509	1393	—	—	85.5 (81.4–89.5)	81.0 (77.0–85.0)	78.8 (74.8–82.7)	75.6 (71.6–79.5)

NH, Non-Hispanic; NHOPI, Native Hawaiian or Other Pacific Islander; —, not applicable.

^a For SUID rates by race and ethnicity, denominator per live births of the specified race and ethnicity.

^b SUID included 3 underlying cause-of-death classifications coded per the ICD-10: SIDS (R95), unknown cause (R99), and ASSB (W75).

^c Race and Hispanic origin are reported separately on birth certificates; persons of Hispanic origin may be of any race. Single race is defined as only 1 race reported on the birth certificate. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Because the 2017 period-linked file was the first year for which all 50 states and Washington DC had birth certificate data on race and Hispanic-origin categories using the 1997 standards, analyses of single race and Hispanic origin groups were restricted to 2017 to 2020.

^d Includes all persons of Hispanic origin of any race.

^e SUID rates with a death count less than 20 are considered “unreliable” and, therefore, not presented.

SIDS and SUID rates from 2019 to 2020 were directly related to infant COVID-19 infection. However, the significant increased rate of SUID among non-Hispanic Black infants from 2019 to 2020, but not among other single race and Hispanic infants, deserves further attention because it could be attributable to the COVID-19 pandemic and its impact on social determinants of health.

The declining SIDS trend during 2015 to 2019 without an increasing overall SUID trend is consistent with trends observed during 1990 to 2015²³ and provides evidence of a continued diagnostic shift. Conversely, the 2020 uptick in the SIDS rate compared to 2019 may be initial evidence of new changes in reporting practices. New guidelines for certification and reporting of

sudden deaths in pediatrics were published by the National Association of Medical Examiners²⁴ and the Radcliffe Congress²⁵ in 2019. If the new guidance was followed, this could have led to increased reporting of SIDS. Both guidance documents state that the presence of sleep environment risk factors (eg, sleep surface sharing, soft bedding, or prone sleep) without adequate evidence for airway obstruction is considered insufficient to certify a death as ASSB and that these deaths may be more appropriately certified as unexplained sudden death or SIDS. However, widespread uptake of this new guidance is unlikely in 1 year’s time, and thus too early, to find direct evidence of an effect of these guidelines on reporting. Jurisdictional changes in reporting would likely occur before nationwide changes are

detectable. If some larger jurisdictions adopted this guidance, it may have affected reporting practices and resulted in the increased US SIDS rate in 2020. Because SUID occurs infrequently, and the number of yearly jurisdictional deaths is small, yearly cause-specific SUID rates by jurisdiction or state could not be explored. As additional years of data become available, it will be important to use the more stable estimates attainable through combined years of data to monitor jurisdictional trends.²³ It is also possible that changes in reporting may have resulted from medical examiner and coroner offices being overburdened by pandemic-related deaths.²⁶ Changes in reporting may have affected the ability to conduct thorough and timely death investigations and autopsies and, consequently, made differentiating




FIGURE 1

SUID and cause-specific SUID rates per 100 000 live births, United States, 2015 to 2020. SUID included 3 underlying cause-of-death classifications coded per the ICD-10: SIDS (R95), unknown cause (R99), and ASSB (W75).

SUID-specific causes more challenging.

Our finding of significantly increased rates of SUID from 2019 to 2020 among non-Hispanic Black infants, but not for other single race or Hispanic infant groups, was unexpected. The COVID-19 pandemic disproportionately affected communities of color compared to non-Hispanic White families through exacerbation of crowded housing, food insecurity,²⁷ unemployment,^{28,29} limited health care access,³⁰ altered child care, and social or emotional health stressors.^{31,32} Infants from these communities were also vulnerable to these effects. Factors exacerbated by the COVID-19 pandemic and

related stay-at-home orders (eg, access to health care, changes to or loss of childcare, parental or caregiver financial and emotional stressors, and increased parental or caregiver substance use) may have also influenced the increased SUID rate among non-Hispanic Black infants in 2020.^{33–35} These factors affect family vulnerability and may have increased unsafe sleep practices (eg, bedsharing), thereby increasing the occurrence of SIDS and other sudden unexpected infant deaths (SUID).^{36,37}

The increasing trend in SUID rates among non-Hispanic Black infants has resulted in a widening racial-ethnic disparity in rates among non-Hispanic Black infants relative to

non-Hispanic White infants (2.8-fold in 2020). The disproportionate burden of COVID-19 illness among communities of color and low-income communities may have widened SUID disparities further, especially among non-Hispanic Black infants.^{38,39} In addition, the persistently higher rates of SUID among non-Hispanic Black and non-Hispanic AIAN infants found in our study and others^{40,41} suggest there may be unique drivers of SUID among these populations, including community-level factors and social determinants of health. Higher rates among these racial and ethnic groups are not unique to SUID; they also exist with other infant health outcomes.⁴² By understanding how social determinants of health




FIGURE 2

Percent distribution of sudden unexpected infant deaths by cause of death and year, United States, 2015 to 2020. Percentages may not sum to 100 because of rounding. SUID included 3 underlying cause-of-death classifications coded per the ICD-10: SIDS (R95), unknown cause (R99), and ASSB (W75).

disproportionately affect infant health and the risks for SUID, health care providers can better support amelioration of those factors.⁴³

Increased clinical training and quality improvement efforts for safe sleep education, and identification of methods for removing barriers to implementing safe sleep practices may be effective in reducing risk, particularly among disproportionately affected populations.⁴⁴ It is important for clinicians to assess additional ways to support, follow-up, and follow-through with families.⁴⁴

Multidisciplinary fatality reviews, such as child death review, and surveillance systems, such as the SUID and Sudden Death in the Young Case Registry, are especially

valuable in increasing understanding of the drivers of the racial disparity and SUID.^{45,46} Multidisciplinary review teams have access to jurisdictional information not available in death certificate data (eg, reports from death scene investigation, autopsy, child protective services, social services, law enforcement, and medical records).⁴⁷ In addition, review teams often have participation from medical providers and social service agencies that know and work directly with families who are affected, and as such, teams can comprehensively review SUID cases, including those that occurred during the COVID-19 pandemic.⁴⁷ With this additional information and active participation from multidisciplinary

professional membership, child death review and the SUID and Sudden Death in the Young Case Registry are well positioned to analyze these data to better understand disparities and risk factors, as well as establish partnerships to drive prevention activities.

A strength and unique quality of our analysis was the application of the single race (as opposed to bridged race) and Hispanic origin groups standards. As of 2020, the bridged race groups that did not follow 1997 OMB guidance are no longer available in birth certificate data, including the linked birth and infant death files. For consistency, the single race standards will need to be




FIGURE 3

Sudden unexpected infant deaths rates per 100 000 live births under 1 year by race and ethnicity, United States, 2017 to 2020. For groups with <100 deaths, statistical significance was assessed by comparing 95% confidence intervals. If confidence intervals overlap, the difference was not statistically significant (NS) at the 95% level.

used to examine SUID rates by race-ethnicity moving forward. Consistent and accurate reporting of race or ethnicity and SUID rate estimates is important for planning the delivery of equitable health services and prioritization of interventions to those infants at highest risk.⁴⁸

This study has a few limitations. First, our analysis could not definitively answer whether SUID rates changed because of the COVID-19 pandemic. With additional years of data, an interrupted time series or Joinpoint analysis will be better able to detect if a changing trend in SUID rates occurred during the COVID-19

pandemic compared to prepandemic years.^{22,49,50} However, this analysis was a valuable first step; early monitoring and continued surveillance activities can identify changes in risk. Second, because our study only included the first year of the pandemic, data were not inclusive of the 2021 Delta and Omicron waves, during which COVID-19 cases (and deaths) were higher among children compared with 2020.⁵¹ Without more years of data, the direct effects of COVID-19 infection on SUID rates could not be determined. However, infants generally experience mild COVID-19 illness, although severe disease and

death do occur in this age group.⁵² Nevertheless, infant mortality declined to a record low in 2020¹ and COVID-19 illness did not appear to contribute to excess infant deaths in 2020 (35 infant deaths and <10 of the SUID had a COVID-19 code). A final limitation is that we did not further disaggregate racial or ethnicity groups. By not disaggregating these groups, we may have masked higher rates of infant mortality, for example among Hispanic subgroups, such as Puerto Rican infants.^{53,54}

We provide new insights about why SIDS rates, as previously reported,¹

may have increased from 2019 to 2020 and present context about other SUID causes (ie, R95, R99, and W75) implicated in reporting variations and subsequent diagnostic shifting.^{11,12} Our findings support evidence that the increased SIDS rate in 2020 as compared to 2019 was likely unrelated to direct effects of the COVID-19 illness but may be attributed to diagnostic shifting in cause-specific SUID rates. Although an overall increased SUID trend during 2015 to 2019 was not observed, continued monitoring is critical; we do not yet understand how the Delta and Omicron COVID-19 variants impacted SUID. Monitoring of indirect factors from the COVID-19 pandemic that may increase the prevalence of SUID risk factors (eg, unsafe sleep environment, smoking, substance use), especially among

racial and ethnic subgroups is also important.⁷ Further investigation and interventions are needed to address the persistent disparity in SUID rates among non-Hispanic Black and non-Hispanic AIAN infants compared to non-Hispanic White infants.

ABBREVIATIONS

AIAN:	American Indian or Alaskan Native
ASSB:	accidental suffocation and strangulation in bed
COVID-19:	coronavirus disease 2019
ICD-10:	International Classification of Diseases, Tenth Revision
OMB:	Office of Management and Budget
SIDS:	sudden infant death syndrome
SUID:	sudden unexpected infant death
WONDER:	Wide-ranging Online Data for Epidemiologic Research

FUNDING: Ms Erck Lambert was supported by a contract between CyberData Technologies, Inc. and the Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, US Centers for Disease Control and Prevention.

CONFLICT OF INTEREST DISCLOSURES: The authors have no potential conflicts of interest relevant to this article to disclose.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

REFERENCES

- Murphy SL, Kochanek KD, Xu J, Arias E. Mortality in the United States, 2020. *NCHS Data Brief*. 2021; (427):1–8
- U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. Healthy People 2030. Social determinants of health. Available at: <https://health.gov/healthypeople/objectives-and-data/social-determinants-health>. Accessed March 16, 2022
- Santoli JM, Lindley MC, DeSilva MB, et al. Effects of the COVID-19 pandemic on routine pediatric vaccine ordering and administration - United States, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(19):591–593
- Ackerson BK, Sy LS, Glenn SC, et al. Pediatric vaccination during the COVID-19 pandemic. *Pediatrics*. 2021;148(1):e2020047092
- Bright Futures Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th ed. American Academy of Pediatrics; 2017
- Mitchell EA, Thach BT, Thompson JM, Williams S. Changing infants' sleep position increases risk of sudden infant death syndrome. New Zealand cot death study. *Arch Pediatr Adolesc Med*. 1999; 153(11):1136–1141
- Moon RY; Task Force on Sudden Infant Death Syndrome. SIDS and other sleep-related infant deaths: evidence base for 2016 updated recommendations for a safe infant sleeping environment. *Pediatrics*. 2016;138(5):e20162940
- Matthews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period linked birth/infant death data set. *Natl Vital Stat Rep*. 2015;64(9):1–30
- Nashelsky MB, Pinckard JK. The death of SIDS. *Acad Forensic Pathol*. 2011; 1(1):92–99
- Shapiro-Mendoza CK, Parks SE, Brustum J, et al. Variations in cause-of-death determination for sudden unexpected infant deaths. *Pediatrics*. 2017; 140(1):e20170087
- Shapiro-Mendoza CK, Tomashek KM, Anderson RN, Wingo J. Recent national trends in sudden, unexpected infant deaths: more evidence supporting a change in classification or reporting. *Am J Epidemiol*. 2006; 163(8):762–769
- Malloy MH, MacDorman M. Changes in the classification of sudden unexpected infant deaths: United States, 1992–2001. *Pediatrics*. 2005;115(5):1247–1253
- Shapiro-Mendoza CK, Palusci VJ, Hoffman B, et al; AAP Task Force on Sudden Infant Death Syndrome, Council on Child

- Abuse and Neglect, Council on Injury, Violence, and Poison Prevention, Section on Child Death Review and Prevention, National Association of Medical Examiners. Half century since SIDS: a reappraisal of terminology. *Pediatrics*. 2021;148(4):e2021053746
14. National Center for Health Statistics. Period linked birth-infant death data files, 2015-2020. Available at: https://www.cdc.gov/nchs/data_access/vitalstatsonline.htm#Period_Linked. Accessed October 7, 2022
15. National Center for Health Statistics. User guide to the 2020 period/2019 cohort linked birth/infant death public use file. Available at: https://ftp.cdc.gov/pub/Health_Statistics/NCHs/Dataset_Documentation/DVS/period-cohort-linked/20PE19CO_linkedUG.pdf. Accessed October 10, 2022
16. National Center for Health Statistics, National Vital Statistics System. Instructions for classifying the underlying cause of death. NCHS Instruction Manual; part 2a. Hyattsville, MD. Available at: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm. Accessed February 22, 2022
17. National Center for Health Statistics, National Vital Statistics System. Instructions for classifying multiple causes of death. NCHS Instruction Manual; part 2b. Hyattsville, MD. Available at: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm. Accessed March 3, 2022
18. World Health Organization. *International Statistical Classification of Diseases and Related Health Problems*. Geneva, Switzerland: World Health Organization; 1992
19. Office of Management and Budget. Revisions to the standards for the classification of Federal data on race and ethnicity. Federal Register 62FR58781-58790, October 30, 1997. Available at: www.whitehouse.gov/wh/eop/omb/html/fedreg.html. Accessed October 8, 2022
20. Schwartz S; National Vital Statistics System. COVID-19 Alert No. 2. New ICD code introduced for COVID-19 deaths. Available at: <https://www.cdc.gov/nchs/data/nvss/coronavirus/Alert-2-New-ICD-code-introduced-for-COVID-19-deaths.pdf>. Accessed February 27, 2022
21. Xu J, Murphy SL, Kochanek KD, Arias E. Deaths: final data for 2019. National Vital Statistics Reports. Available at: <https://stacks.cdc.gov/view/cdc/106058>. Accessed October 17, 2022
22. Ingram DD, Malec DJ, Makuc DM, et al. National Center for Health Statistics guidelines for analysis of trends. *Vital Health Stat 2*. 2018; (179):1–71
23. Erck Lambert AB, Parks SE, Shapiro-Mendoza CK. National and state trends in sudden unexpected infant death: 1990-2015. *Pediatrics*. 2018;141(3):e20173519
24. Bundock EA, Eason EA, Andrew TA, et al. Death certification and surveillance. In: Bundock EA, Corey TS, Andrew TA, et al, eds. *Unexplained Pediatric Deaths: Investigation*. San Diego, CA: Certification, and Family Needs; 2019
25. Goldstein RD, Blair PS, Sens MA, et al; 3rd International Congress on Sudden Infant and Child Death. Inconsistent classification of unexplained sudden deaths in infants and children hinders surveillance, prevention and research: recommendations from The 3rd International Congress on Sudden Infant and Child Death. *Forensic Sci Med Pathol*. 2019;15(4):622–628
26. Moretti M, Malhotra A, Visonà SD, Finley SJ, Osculati AMM, Javan GT. The roles of medical examiners in the COVID-19 era: a comparison between the United States and Italy. *Forensic Sci Med Pathol*. 2021;17(2):262–270
27. Cassidy-Vu L, Way V, Spangler J. The correlation between food insecurity and infant mortality in North Carolina. *Public Health Nutr*. 2022;25(4):1038–1044
28. Couch KA, Fairlie RW, Xu H. Early evidence of the impacts of COVID-19 on minority unemployment. *J Public Econ*. 2020;192:104287
29. Monte LM, Perez-Lopez DJ. America counts: stories behind the numbers. How the pandemic affected Black and White households. Available at: <https://www.census.gov/library/stories/2021/07/how-pandemic-affected-black-and-white-households.html>. Accessed March 21, 2022
30. Hambidge SJ, Emsermann CB, Federico S, Steiner JF. Disparities in pediatric preventive care in the United States, 1993-2002. *Arch Pediatr Adolesc Med*. 2007;161(1):30–36
31. McKnight-Eily LR, Okoro CA, Strine TW, et al. Racial and ethnic disparities in the prevalence of stress and worry, mental health conditions, and increased substance use among adults during the COVID-19 pandemic - United States, April and May 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(5):162–166
32. Magesh S, John D, Li WT, et al. Disparities in COVID-19 outcomes by race, ethnicity, and socioeconomic status: a systematic-review and meta-analysis. *JAMA Netw Open*. 2021;4(11):e2134147
33. Patrick SW, Henkhaus LE, Zickafoose JS, et al. Well-being of parents and children during the COVID-19 pandemic: a national survey. *Pediatrics*. 2020;146(4):e2020016824
34. Czeisler ME, Lane RI, Wiley JF, Czeisler CA, Howard ME, Rajaratnam SMW. Follow-up survey of US adult reports of mental health, substance use, and suicidal ideation during the COVID-19 pandemic, September 2020. *JAMA Netw Open*. 2021;4(2):e2037665
35. Lebrun-Harris LA, Ghandour RM, Kogan MD, Warren MD. Five-year trends in US children's health and well-being, 2016-2020. *JAMA Pediatr*. 2022;176(7):e220056
36. Shipstone RA, Young J, Kearney L, Thompson JMD. Applying a social exclusion framework to explore the relationship between sudden unexpected deaths in infancy (SUDI) and social vulnerability. *Front Public Health*. 2020;8:563573
37. Wood AM, Pasupathy D, Pell JP, Fleming M, Smith GC. Trends in socioeconomic inequalities in risk of sudden infant death syndrome, other causes of infant mortality, and stillbirth in Scotland: population based study. *BMJ*. 2012;344:e1552
38. Parcha V, Booker KS, Kalra R, et al. A retrospective cohort study of 12,306 pediatric COVID-19 patients in the United States. *Sci Rep*. 2021;11(1):10231
39. Figueroa JF, Wadhera RK, Lee D, Yeh RW, Sommers BD. Community-level factors associated with racial and ethnic disparities in COVID-19 rates in Massachusetts. *Health Aff (Millwood)*. 2020;39(11):1984–1992

40. Devin J, Marano R, Mikhael M, Feaster W, Sanger T, Ehwerhemuepha L. Epidemiology of neonatal COVID-19 in the United States. *Pediatrics*. 2022;150(4): e2022056297
41. Parks SE, Erck Lambert AB, Shapiro-Mendoza CK. Racial and ethnic trends in sudden unexpected infant deaths: United States, 1995-2013. *Pediatrics*. 2017;139(6):e20163844
42. MacDorman MF. Race and ethnic disparities in fetal mortality, preterm birth, and infant mortality in the United States: an overview. *Semin Perinatol*. 2011;35(4):200–208
43. Hwang SS, Melvin P, Diop H, Settle M, Mourad J, Gupta M. Implementation of safe sleep practices in Massachusetts NICUs: a state-wide QI collaborative. *J Perinatol*. 2018;38(5): 593–599
44. Horbar JD, Edwards EM, Ogbolu Y. Our responsibility to follow through for NICU infants and their families. *Pediatrics*. 2020;146(6):e20200360
45. Shapiro-Mendoza CK, Camperlengo LT, Kim SY, Covington T. The sudden unexpected infant death case registry: a method to improve surveillance. *Pediatrics*. 2012;129(2):e486–e493
46. Covington TM. The US National Child Death review case reporting system. *Inj Prev*. 2011;17(Suppl 1):i34–i37
47. The National Center for Fatality Review and Prevention. *Keeping Kids Alive: Child Death Review in the United States, 2020*. Okemos, MI: Michigan Public Health Institute; 2021
48. Yoon P, Hall J, Fuld J, et al. Alternative methods for grouping race and ethnicity to monitor COVID-19 outcomes and vaccination coverage. *MMWR Morb Mortal Wkly Rep*. 2021;70(32):1075–1080
49. Penfold RB, Zhang F. Use of interrupted time series analysis in evaluating health care quality improvements. *Acad Pediatr*. 2013;13(6 Suppl):S38–S44
50. Bernal JL, Cummins S, Gasparrini A. Interrupted time series regression for the evaluation of public health interventions: a tutorial. *Int J Epidemiol*. 2017; 46(1):348–355
51. Marks KJ, Whitaker M, Anglin O, et al; COVID-NET Surveillance Team. Hospitalizations of children and adolescents with laboratory-confirmed COVID-19 - COVID-NET, 14 states, July 2021-January 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71(7):271–278
52. Ouldali N, Yang DD, Madhi F, et al; investigator group of the PANDOR study. Factors associated with severe SARS-CoV-2 infection. *Pediatrics*. 2021;147(3): e2020023432
53. Ely DM, Driscoll AK. Infant mortality in the United States, 2018: data from the period linked birth/infant death file. *Natl Vital Stat Rep*. 2020;69(7):1–18
54. Office of Minority Health, U.S. Department of Health and Human Services. *Infant Mortality and Native Hawaiians/Pacific Islanders*. Available at: <https://minorityhealth.hhs.gov/omh/browse.aspx?lvl=4&lvlid=83>. Accessed April 19, 2022